

# Smokeless Tobacco Use (Adults)

## Smokeless Tobacco Use (Adults)

- Most recent data (2013): 3% or about 38,300 King County adults age 18+.
- Trend over time: 2000-13: flat
- Description: Currently use chewing tobacco, snuff, or snus

## Smokeless tobacco use (adults), King County, 2009-2013 average

	Percent	Lower CI	Upper CI
<b>King County</b> (adults age 18+)	3	2	3
<b>AGE</b>			
18-24	5	3	7
25-44	4	3	5
45-64	2	1	2
65+	1	0	1
<b>GENDER</b>			
Male	5	4	6
Female	1	0	1
<b>RACE/ETHNICITY</b>			
AIAN	2 §	1 §	6 §
Asian	3 §	2 §	6 §
Black	3 §	1 §	6 §
Hispanic	3 §	1 §	5 §
Multiple	3 §	1 §	9 §
NHPI	9 §	3 §	27 §
White	3	2	3
<b>HOUSEHOLD INCOME</b>			
<\$15,000	5 §	3 §	9 §
\$15,000 to \$24,999	4	2	6
\$25,000 to \$34,999	2 §	1 §	5 §
\$35,000 to \$49,999	3	2	5
\$50,000 to \$74,999	3	2	5
\$75,000+	2	1	3
<b>REGION</b>			
East	3	2	4
North	2 §	1 §	5 §
Seattle	2	1	3
South	3	2	4

### Comparisons:

Higher than King County rate (yellow)

Lower than King County rate (green)

### Notes:

Source: Behavioral Risk Factor Surveillance System.

Prepared by Public Health - Seattle & King County, APDE, 01/2015.

CI is 95% Confidence Interval.


\* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

## Smokeless tobacco use (adults)

### King County, 2009-2013 average



Source: Behavioral Risk Factor Surveillance System.

Prepared by Public Health - Seattle & King County, APDE, 01/2015.

[-----] Confidence interval shows range that includes true value 95% of the time.

\* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.

## Smokeless tobacco use (adults) by comparison areas

The Healthy People 2020 Objective for smokeless tobacco use (adults) is 0%.

	Latest Available Year	Percent	Lower CI	Upper CI
United States	2010	3	2	3
WA State	2012	4	3	4
<b>King County</b>	2013	3	2	3


### Notes:

Source: Behavioral Risk Factor Surveillance System.

Prepared by Public Health - Seattle & King County, APDE, 01/2015.

CI is 95% Confidence Interval.

## Smokeless tobacco use (adults) by comparison areas



Source: Behavioral Risk Factor Surveillance System.  
Prepared by Public Health - Seattle & King County, APDE, 01/2015.  
|-----| Confidence interval shows range that includes true value 95% of the time.

# Smokeless tobacco use (adults) by health reporting areas, King County, 2009-2013 average

	Percent	Lower CI	Upper CI
<b>King County</b> (adults age 18+)	3	2	3
Auburn	3 §	1 §	8 §
..Auburn-North	2 §	0 §	10 §
..Auburn-South	5 §	1 §	14 §
Bear Creek/Carnation/Duvall	5 §	2 §	11 §
Bellevue	2 §	1 §	5 §
..Bellevue-Central	3 §	1 §	8 §
..Bellevue-NE	4 §	1 §	12 §
..Bellevue-South	1 §	0 §	5 §
..Bellevue-West	1 §	0 §	5 §
Black Diamond/Enumclaw/SE County	4 §	1 §	10 §
Bothell/Woodinville	1 §	0 §	3 §
Burien	1 §	0 §	4 §
Covington/Maple Valley	5 §	2 §	9 §
Des Moines/Normandy Park	4 §	1 §	9 §
East Federal Way	9 §	3 §	24 §
Fairwood	2 §	0 §	7 §
Federal Way	3 §	1 §	8 §
..Fed Way-Central/Military Rd	5 §	2 §	12 §
..Fed Way-Dash Point/Woodmont	1 §	0 §	3 §
Issaquah	4 §	2 §	10 §
Kenmore/LFP	2 §	0 §	6 §
Kent	2 §	1 §	5 §
..Kent-East	4 §	1 §	12 §
..Kent-SE	1 §	0 §	3 §
..Kent-West	5 §	1 §	18 §
Kirkland	2 §	1 §	4 §
..Kirkland	1 §	0 §	4 §
..Kirkland North	2 §	1 §	7 §
Mercer Isle/Pt Cities	0 §	0 §	2 §
Newcastle/Four Creeks	1 §	0 §	3 §
North Highline	0 §	0 §	3 §
Redmond	4 §	1 §	13 §
Renton	1 §	0 §	3 §
..Renton-East	1 §	0 §	5 §
..Renton-North	1 §	0 §	5 §
..Renton-South	1 §	0 §	5 §
Sammamish	2 §	1 §	5 §
SeaTac/Tukwila	9 §	4 §	18 §
Seattle	2	2	3
..Ballard	2 §	1 §	4 §
..Beacon/Gtown/S.Park	5 §	2 §	13 §
..Capitol Hill/E.lake	1 §	0 §	3 §
..Central Seattle	4 §	1 §	11 §
..Delridge	3 §	1 §	7 §
..Downtown	1 §	0 §	5 §
..Fremont/Greenlake	2 §	0 §	7 §
..NE Seattle	2 §	1 §	6 §
..North Seattle	0 §	0 §	1 §
..NW Seattle	2 §	1 §	7 §
..QA/Magnolia	3 §	1 §	6 §
..SE Seattle	1 §	0 §	5 §
..West Seattle	1 §	0 §	3 §
Shoreline	1 §	0 §	3 §
Snoqualmie/North Bend/Skykomish	2 §	1 §	5 §
Vashon Island	2 §	1 §	9 §

## Comparisons:

- Higher than King County rate (yellow)
- Lower than King County rate (green)

## Notes:

Source: Behavioral Risk Factor Surveillance System.

Prepared by Public Health - Seattle & King County, APDE, 01/2015.

CI is 95% Confidence Interval.


\* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.

Persons of Hispanic ethnicity can be of any race and are included in the racial categories.




## Smokeless tobacco use (adults) by health reporting areas King County, 2009-2013 average



Note: HRA labels on the map match the chart on the right, listed in alphabetical order  
Source: Behavioral Risk Factor Surveillance System.  
Prepared by Public Health - Seattle & King County, APDE, 01/2015.

# Smokeless tobacco use (adults) King County, 2000-2013, three-year rolling averages



Source: Behavioral Risk Factor Surveillance System.  
Prepared by Public Health - Seattle & King County, APDE, 01/2015.

Produced by the  
Assessment, Policy Development & Evaluation Unit  
Public Health - Seattle & King County  
01/15

King County  
Hospitals  
for a Healthier  
Community